

Boise State University
ScholarWorks

College of Arts and Sciences Poster Presentations

2010 Undergraduate Research and Scholarship
Conference

4-12-2010

Population Dynamics of Ten Avian Species Breeding at Lucky Peak, Boise Foothills, Idaho

Heidi Ware

Department of Biological Sciences, Boise State University

2nd Place Winner of the 2010 Student Choice Award.



Population Dynamics of Ten Avian Species Breeding at Lucky Peak, Boise Foothills, Idaho

Heidi Ware, Dr. Jay Carlisle, & Dr. Alfred Dufty
Department of Biological Sciences, Boise State University, Boise, Idaho, 83725, USA



ABSTRACT

We explored capture and banding data from 1997-2009 at the Idaho Bird Observatory's Lucky Peak banding station to examine patterns of abundance and survivorship for the ten most frequently-captured breeding bird species, including how these patterns varied by year. We found that survivorship patterns vary by species. We documented Nashville and Yellow warblers surviving up to four years whereas multiple individuals of Dusky Flycatcher, MacGillivray's Warbler, and Chipping Sparrow lived seven years or more. In fact, several Western Tanagers and Dusky Flycatchers exceed or match longevity records according to the USGS Bird Banding Lab's database.

The ratio of hatch-year birds (0 years old) to adult birds (1+ year old) lends insight into survivorship, productivity, and recruitment within Lucky Peak's breeding bird community. The ratios observed in Warbling Vireos and other species suggests that immigration is important for maintaining their breeding populations at Lucky Peak. In contrast, ratios for other species implied that many individuals that are hatched at the site return to breed.

Our analysis also revealed trends in population dynamics and striking annual differences in species' abundance. Neotropical migrant warblers such as MacGillivray's Warbler and Yellow Warbler peaked in both 2001 and 2007. Interestingly, short-distance migrant sparrows such as Chipping Sparrow and Spotted Towhee peaked in 2004.

Preliminary investigations into weather trends showed some meaningful correlations for the species analyzed. Precipitation and temperatures from Boise, ID (breeding grounds) and Guadalajara, Mexico (approximate wintering grounds for Neotropical migrants) were correlated to yearly population changes for some species. MacGillivray's Warbler populations are known to winter in Mexico near Guadalajara and were influenced by winter precipitation whereas Audubon's Warbler and Warbling Vireo year to year numbers were influenced more by springtime temperature and precipitation in Boise.



INTRODUCTION

Goals of study:

- Explore variables that effect populations of passerine bird species breeding at Lucky Peak near Boise, ID

Many variables cause animal populations to fluctuate annually, including:

- habitat change
- changes in predator-prey interactions
- weather
- disease

This study explored population changes caused by:

- life history traits
 - diet – seeds versus insects
 - migratory strategies– distance, location, timing
- weather
 - breeding grounds (Boise)
 - wintering grounds (Guadalajara)

Use recapture data to determine longevity

- Analyze weather data from breeding (Boise) and wintering grounds (Guadalajara) to determine its affects on population
- Compare differences in species' diet and migratory strategy to explain population trends.
- Use age ratios of Hatch Year (0 year old) and Second Year (1 year old) birds to infer survivorship and recruitment

Hypotheses:

- Species with few young produced locally must rely on immigration to maintain populations
- Species with different diets or migration strategies will differ in their patterns of variation in annual success
- Weather may be a significant cause of the observed population fluctuations

METHODS

- Utilized 13 years of data from Lucky Peak for the 10 most common breeding species
- Calculated minimum possible age for longevity data
- Population abundance calculated
- Regression analysis used to correlate population and weather trends for 2000-2009
- airport temperature and precipitation data used for weather analysis
 - regression analysis of Boise data performed for:
 - Audubon's Warblers, Warbling Vireos, Dusky Flycatchers
 - analysis of Guadalajara data done for:
 - MacGillivray's Warblers, Warbling Vireos, Dusky Flycatchers

RESULTS

Species listed in order of abundance

(n=total individuals captured during the 2000-2009 breeding seasons)

- Dusky Flycatcher (n=1,805)
- MacGillivray's Warbler (n=1,736)
- Chipping Sparrow (n=1,115)
- Yellow Warbler (n=1,035)
- Nashville Warbler (n=1,006)
- Audubon's Warbler (n=976)
- Spotted Towhee (n=763)
- Warbling Vireo (n=713)
- Western Tanager (n=625)
- Lazuli Bunting (n=564)

Age ratios

- Many second year (SY; 1 yr old) 'adult' birds and few locally-produced hatch year (HY; 0 yr old) birds in populations of:
 - Warbling Vireo (Figure 1)
 - Western Tanager
 - Dusky Flycatcher
- Other species populations had equal proportions of HY & SY birds

Life Histories (Figure 2)






- Chipping Sparrows and Spotted Towhees:
 - populations peak near 2004
- Yellow and MacGillivray's Warblers:
 - populations peak near 2001 and 2007
 - populations decrease in 2004

Longevity

- Maximum age for individuals:
 - range: 4 to 10 years
 - most species: 6 or 7 years
- Several species matched or exceeded age records:
 - Dusky Flycatcher:
 - BBL (Bird Banding Lab) record: 8 years
 - Lucky Peak records: one age 10, one age 9
 - Western Tanager:
 - BBL and Lucky Peak record: 6 years.
- Some species were well below BBL records:
 - Yellow and Nashville Warblers:
 - BBL record: 10 years
 - Lucky Peak record: 4 years

Weather

- Boise weather has significant impacts on populations.
- Springtime temperatures and precipitation influenced:
 - Audubon's Warblers (Figure 3)
 - greater June rainfall = higher #s of HY birds
 - Warbling Vireos (Figure 4)
 - warmer average spring temp = higher numbers of total birds
- Guadalajara weather had impacts on MacGillivray's Warbler populations (Figure 5).
- more February precipitation = higher numbers of MacGillivray's Warblers



PRELIMINARY CONCLUSIONS

Age ratios (Figure 1)

- Few HY birds in one year and more SY birds the following year strongly suggests:
 - not all SY birds were hatched at Lucky Peak
 - recruitment (immigration of new individuals into a population) is important for these species
 - adult population is not sustained by young produced at the site
- Other species show fewer SY birds relative to HY's the following year which suggests:
 - the population does not have high recruitment
 - many birds hatched at Lucky Peak likely return as adults

Life Histories

- Chipping Sparrows and Spotted Towhees are similar:
 - in the sparrow family *Emberizidae*
 - rely primarily on seeds in winter
 - winter in Texas and N. Mexico (Figure 6)
- MacGillivray's and Yellow Warblers are similar:
 - in the warbler family *Parulidae*
 - feed primarily on insects throughout the year
 - Neotropical migrants: winter in W. & S. Mexico (Figure 7)

Diet Hypothesis

- 2004: good year for seeds?
- 2001 and 2007: good years for insects?

Migration Hypothesis

- arrival and breeding times differ because of distance traveled
- wintering locations or migration routes had different:
 - weather
 - habitat changes
 - food supplies

True cause is unknown!

Longevity

Multiple species studied did not match the BBL longevity data

- Dusky Flycatchers exceeded BBL records:

- Why?
 - more multi-year recaptures. Increased odds because:
 - Dusky's may survive longer at Lucky Peak *and/or*
 - higher site fidelity here than at other locations
- Yellow and Nashville Warblers were below BBL records
- Why?
 - Lucky Peak breeders have lower survivorship *and/or*
 - not enough recaptures to show true age distribution

Weather

Warmer temps and more rainfall in Boise increased population numbers for all species examined.

- Springtime temps (Figure 4) may influence:
 - arrival timing of adults
 - nesting success
 - food supplies

- Rainfall in June (Figure 3) could influence:
 - amount of food available to young birds

- June is an important month for nesting and fledging
- More rainfall in Guadalajara increased numbers of MacGillivray's Warblers.
- Correlation supports idea that Guadalajara is near actual wintering grounds (or an area with similar climate) for the Lucky Peak population (Figure 5)
- February precipitation may mean:
 - good weather in general
 - improved food supplies:
 - increased fitness of adults prior to migration
 - more likely to survive migration journey
 - more fit for reproduction once they arrive

Similar conclusions were also reached by Nott et al. 2002!

Special thanks to:

- Rob Miller for invaluable statistics advice and computer help
- The Idaho Bird Observatory and all crews from '97-'09 who contributed to the data set

